

Review

Molecular makeup of the human adult ovary

 Xueying Fan¹ and Susana M. Chuva de Sousa Lopes^{1,2}

Abstract

Functional ovarian cells are essential for human fertility. In the adult ovary, different cell types ensure ovary homeostasis, enable hormonal production, and support oocyte maturation. Hence, the ovary is a complex and highly dynamic organ composed of a great diversity of cell types, with many still uncharacterized. The use of single-cell RNA sequencing technologies on human ovarian tissue is starting to unravel the molecular signature of the cells present in the ovary, highlighting dramatic changes in gene expression during follicular growth and regression. This knowledge will ultimately provide insights into female fertility and associated reproductive diseases and will allow the optimization of human-based disease models and in vitro gametogenesis protocols.

Addresses

¹ Department of Anatomy and Embryology, Leiden University Medical Center, Einthovenweg 20, 2333 ZC Leiden, the Netherlands

² Department for Reproductive Medicine, Ghent University Hospital, Corneel Heymanslaan 10, 9000 Ghent, Belgium

Corresponding author: Chuva de Sousa Lopes, Susana M. (lopes@lumc.nl)

Current Opinion in Endocrine and Metabolic Research 2021, 18:187–193

This review comes from a themed issue on Regulation of Ovarian Function

Edited by Norah Spears and Kate Hardy

For complete overview of the section, please refer the article collection - Regulation of Ovarian Function

Available online 8 April 2021

<https://doi.org/10.1016/j.coemr.2021.03.016>

2451-9650/© 2021 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Keywords

Human, Adult ovary, Transcriptional signature, Single-cell sequencing, Folliculogenesis.

Abbreviations

TC, theca cell; GC, granulosa cell; mm, millimeter; GV, germinal vesicle; GVBD, germinal vesicle breakdown; LH, luteinizing hormone; MI, metaphase I; MII, metaphase II; OSE, ovarian surface epithelium; COC, cumulus–oocyte complex.


Introduction

The human ovary is a highly complex and dynamic reproductive exocrine organ composed of many different cell types (Figure 1a) and responsible for storage and

maturation of oocytes and cyclic production of hormones, such as progesterone and estrogen. The maturation of a primary oocyte into a competent oocyte, ready for ovulation, is a process tightly regulated by ovarian somatic cells [1–3]. Each oocyte is encapsulated by somatic cells, forming a functional unit known as the follicle. The two main somatic cell types responsible for formation and growth of the follicle are granulosa cells (GCs), surrounding the oocyte, and theca cells (TCs), aligned on the outer surface of the basal lamina [4,5]. Those two lineages undergo a dramatic development during follicular growth/maturation, followed by either regression or further development into the corpus luteum (Figure 1b). Depending on the stage of the follicle (or subsequent corpus luteum), the GC morphology can be subdivided into squamous, cuboidal, mural, cumulus, corona radiata, and lutein GCs (Figure 1). The development of the TC lineage is less well understood, but includes several well-known subtypes, such as interna, externa, and lutein TC (Figure 1). As the GC and TC develop, the follicle transits from primordial, primary, secondary (<0.15 mm), preantral (0.15–0.2 mm) (Figures 1b, 2a–d) to antral follicles (0.2–5 mm), when early GCs separate into cumulus GCs surrounding the oocyte and mural GC lining the basal lamina (Figures 1b, 2e–f). In the ovary, there are many antral follicles, but per menstrual cycle, typically one will become dominant and increase in size, reaching about 20 mm before ovulation [6]. After ovulation, that follicle transforms into the corpus luteum containing lutein GCs and lutein TCs (Figure 2g). Inevitably, the corpus luteum undergoes luteolysis and regresses into the corpus albicans [7,8], whereas most antral follicles that undergo atresia degenerate [9], leaving behind the Slavjanski's glassy membrane (basement membrane) as the only remnant (Figure 1b).

Owing to recent developments in single-cell sequencing technology, we are starting to gain in-depth knowledge on the different cell types present in the adult ovary that are responsible for follicular growth and degeneration as well as the extraordinary tissue remodeling that takes place in the human adult ovary during each menstrual cycle [10–13]. In addition, the ovary also contains many other cell types, such as cells of the ovarian surface epithelium (OSE) and stromal cells that include a variety of immune cells, endothelial cells of blood and lymph vessels, smooth muscle cells, and fibroblasts of the tunica albuginea and ovarian medulla [14] (Figure 1).

Figure 1



Main cell types in the human adult ovary. (a) Cartoon overview of the cellular makeup present in the human adult ovary and (b) the development of the oocyte, granulosa cell (GC), and theca cell (TC) during folliculogenesis. GV, germinal vesicle; GVBD, germinal vesicle breakdown; MI, metaphase I; MII, metaphase II.

Molecular dynamics of oocytes during folliculogenesis


Primary oocytes are arrested at dictyate (diplotene of meiotic prophase I), in the so-called germinal vesicle (GV) stage, and are present in the ovarian cortex, where they constitute a finite pool of resting follicles that have not started to grow. After a surge in luteinizing hormone in the preovulatory follicle, meiotic resumption starts with germinal vesicle breakdown, but pauses once more at metaphase II (MII), until it concludes with the extrusion of the second polar body after fertilization [15] (Figure 1b). Several well-known germ cell and oocyte-specific genes, including *DDX4*, *DPPA3*, *GDF9*, *OOSP2*, *ZP2*, *ZP3*, *FIGLA*, *LHX8*, *SOHLH2*, *DAZL*, and *SOX30*, are detected across maturation stages [12,13,16–18] (See Figure 3).

Laser capture microdissection followed by transcriptional analysis has revealed substantial differential expression between human oocytes from primordial and

primary follicles, such as high *FOXO1*/*FOXO1* expression in oocytes in primordial follicles and high *EIF4E*/*EIF4E* expression in oocytes in primary follicles [19]. In addition, nuclear localization of *FOXO3* and *PTEN* is involved in maintaining dormancy, whereas nuclear accumulation of phosphorylated Akt and phosphorylated mTOR initiates the transition from primordial to primary follicles [20]. In agreement, activation of the PI3K/Akt/mTOR pathway stimulates this transition [21]. Single-cell transcriptomics has further revealed that oocytes in primary follicles express high levels of *CLVS1*, *STAT4*, *TLR4*, *CDK14*, and *DDIT4L*, whereas oocytes in secondary follicles are enriched in *USP27X*, *PAEP*, *COX7A1*, and *CHST7*, and oocytes in early antral follicles show upregulation of *NTF4*, *LCP2*, *NSMF*, and *ABCD1* [13] (See Figure 3).

Microarray analysis of preovulatory oocytes from the GV stage to MII showed that expression of *PCNA*, *FGF14*, *BMP15*, and *CDC25A* are largely increased in MII

Figure 2



Images of granulosa cells (GCs) and theca cells (TCs) during folliculogenesis in humans. Histological images showing hematoxylin and eosin staining of (a) primordial follicles, (b) primary follicles, (c) secondary follicles, (d) preantral follicles, and (e) antral follicles, with (f) an insert magnified and (g) the corpus luteum showing different types of GCs and TCs. Scale bar in a–d and f is 10 μ m, in e is 100 μ m, and in g is 50 μ m.

oocytes compared with GV and metaphase I oocytes [22]. More recently, high expression of *RBBP7* was observed in MII oocytes compared with metaphase I oocytes using single-cell transcriptomics [23]. In addition, several studies have used single-cell transcriptomics to compare the oocyte quality of patients of different ages with different reproductive disorders (endometriosis, polycystic ovary syndrome) or oocytes undergoing in vitro maturation with those of healthy females [24–32]; however, owing to the low number of available oocytes, a robust characterization of GV to MII progression is still lacking.

Molecular dynamics of the GC during folliculogenesis


During folliculogenesis, GCs are the only cells that interact directly with oocytes, having a decisive impact on oocyte quality (Figure 2). In primordial follicles (fetal and adult), squamous GCs are known to express *FOXL2*, *GATA6*, *GATA4*, and *WT1* [12,33,34]. Several signaling pathways, such as mTOR, Hippo, and TGF β superfamily, have been associated with the transition from the primordial (squamous GC) to primary (cuboidal/early GC) follicle [21,35–37]. As per RNA sequencing performed in samples of 10 pooled GCs, GCs from the primary follicle were shown to be enriched in *RND3*, *MGP*, and *TST*; GCs from secondary follicles were

enriched in *BNIP1*, whereas levels of several hormone receptors (*AR*, *PGR*, *ESR1*, *FSHR*) and genes related to steroidogenesis (*CYP11A1*, *CYP19A1*, *HSD3B2*, *HSD17B1*) were increased in GCs in antral follicles [13].

Single-cell transcriptomics on 1- to 5-mm antral follicles has revealed that many markers for GCs, such as *AMH*, *CDH2*, *GJA1*, *TNNI3*, *GSTA1*, *CD99*, *SERPINE2*, *DSP*, *MAGED2*, *SPINT2*, and *TSPAN6*, were observed in all GC subtypes [10,11] (See Figure 3). Moreover, the majority of GCs from small antral follicles (1–2 mm) showed high levels of *WT1* and *EGR4*, characteristic of an early GC signature, before differentiation into cumulus and mural GCs. A clear molecular signature separating cumulus and mural GCs was only observed in antral follicles larger than 2 mm, with high levels of *KRT18*, *CITED2*, and *AKIRIN1* observed in mural (and early) GCs and high levels of *IGFBP2*, *HTRA1*, *INHBB*, and *IHH* specifically expressed in cumulus GCs [10] (See Figure 3).

Cumulus GCs collected from cumulus–oocyte complexes from patients undergoing medical assisted reproduction and containing either GV or MII oocytes showed differential expression of cellular matrix components (*VTN*, *SPOCK2*, *TNC*) and genes involved in steroid metabolism and processing (*AREG*, *EREG*, *HSD11B1*,

Figure 3



Highly differentially expressed genes in human adult ovarian cells. Overview of highly differentially expressed genes in main cell types in the ovary (oocytes and somatic cells) in different stages of folliculogenesis. COC, cumulus–oocyte complex; GC, granulosa cell; TC, theca cell.

STAR, DHCR24) with upregulation in cumulus GCs associated with MII oocytes [13,38]. Interestingly, immune and inflammation factors are also found upregulated in cumulus GCs associated with MII oocytes [38,39]. In accordance, cumulus GCs isolated at different time points during ovarian stimulation and subsequent ovulation also showed upregulation of inflammation-related genes, such as *IL7R, CXCR4*, and *CD24* [40]. Moreover, expression of genes involved in cumulus expansion (*TNFAIP6, PTX3, HAS2, FGG*) and genes potentially involved in follicle rupture and tissue remodeling (*TIMP1, ADAMTS1, ADAMTS9*) were shown to be increased until ovulation by transcriptomics on cumulus GCs and/or by proteomics of follicular fluid [40,41].

During the formation of the corpus luteum, mural GCs differentiate into lutein GCs (Figures 1b and 2g). In the corpus luteum, *CYP19A1* and *VEGF* are expressed exclusively in lutein GCs, whereas *HSD3B2, STAR*, and *CYP11A1* are expressed by both lutein GCs and lutein TCs [7,42–44] (See Figure 3). During the collection of cumulus–oocyte complexes, additional mural GCs are also aspirated, and those can be transiently cultured and induced to become lutein GCs in vitro. Using this culture

model, *SCN9A, OTR*, and *PEDF* have been shown to be involved in the regression of lutein GCs [45].

Molecular dynamics of TCs during folliculogenesis

In the ovary, the origin and function of TCs are not well understood, but TCs are known to be involved in diverse aspects of folliculogenesis that include endocrine, structural, vascular, and immune functions [4,46].

Similarly to early GCs in 1- to 2-mm antral follicles, TCs in those follicles show a unique molecular signature that includes expression of *MME2, APOD, APOC1, MEST*, and *WFDC1* [10]. It is only in larger antral follicles that early TCs differentiate into endocrine interna TCs and fibroblast-like externa TCs [47]. In agreement, genes involved in ovarian hormonal biosynthesis and lipid metabolism, such as *STAR, CYP11A1, CYP17A1, LHCGR, FDX1, INSL3, APOA1*, and *PLIN2*, are expressed in the interna TC; by contrast, externa TCs are enriched in genes associated with production of extracellular matrix and vasculature development, such as *COL1A1, COL1A2, COL3A1, COL3A2, COL6A1, FGFR1, STAT3*, and *ACTA2* [10,11,42] (See Figure 3). Interna and externa TCs can be

separated owing to the differential expression of cell surface makers: ANPEP (CD13) is specifically expressed in interna TCs, whereas externa TCs are positive for ENG (CD105) and GPC3 [11]. Lutein TCs sustain the expression of steroidogenic genes, such as CYP17A1 [7,48].

Molecular makeup of the ovarian stroma

Many cell types present in the ovarian stroma can also be found in the stroma of other organs in the body. Those include a variety of immune cells, endothelial cells of lymph and blood vasculature, smooth muscle cells, and fibroblast-like cells. Some of those cell lineages may express organ-specific genes, but they also express canonical lineage markers, such as *PTPRC*, *CD53*, and *CXCR4* in immune cells; *VWF*, *CLDN5*, *CD34*, and *CDH5* in endothelial cells; *TAGLN*, *MYH11*, and *RG55* in smooth muscle cells; and *DCN*, *TCF21*, and *ARID5B* in fibroblast-like cells [10–12] (See Figure 3). Nevertheless, it is worth noting that each main lineage is heterogeneous, containing several cellular subtypes. For example, the ovary contains several types of immune cells such as monocytes, macrophages, B cells, T cells, and natural killer (NK) cells [10,12], and most probably, those sub types can be further subdivided into additional subtypes. In this regard, also, different subtypes of fibroblast-like cells, perhaps present in different parts of the ovary (tunica albuginea, inner cortex, and medulla), different types of epithelial cells, and smooth muscle cells are responsible for supporting the dynamic structure of the ovary [14]. During the cyclic transition between the follicular and luteal phase, different cell types ensure follicular growth and remodeling, contributing to the great temporal and spatial cellular complexity observed in the ovary.

One of the cell populations that have not yet been captured using single-cell sequencing technologies is the OSE, a single layer of mesothelium cells that covers the ovaries, interfacing with the peritoneal cavity. The role of the OSE is largely obscure. During fetal development, it may contribute to form GCs, whereas in the adult, it may contribute to repair the ovarian surface after ovulation [49]. Regarding this repair function, cells of the OSE express several markers associated with adult stem cells, such as *PROM1* (*CD133*), *RSP01*, *LGR5*, *CD44*, *ALDH1A1*, and several types of keratins (*KRT18*, *KRT19*), but are negative for *MUC16* (*CA125*) [50] (See Figure 3). The functional relationship between the mesothelium cells of the OSE and the mesothelium cells of the fallopian tubes (fimbria) remains obscure, but they both have been associated with a common type of epithelial ovarian cancer, high-grade serous ovarian carcinoma [51].

Summary

The transcriptional signature of oocytes, GCs, and TCs among all other cells present in the adult ovary is

starting to reveal the cellular complexity and dynamics of the human ovary. In future, including more samples of different phases of the ovarian cycle and perhaps ovarian tissue from patients with different ages and reproductive diseases will allow a more complete understanding of the broad range of cell types in the ovary, which will benefit the development of disease models. Moreover, the differentiation in vitro of human competent oocytes, starting from primordial follicles, is still a challenge. To mimic follicle development in vitro, the molecular analysis of ovarian cells, including oocytes, GCs, and TCs, during culture will allow efficient optimization of culture protocols to achieve oocyte maturation, providing innovative solutions for medical assisted reproduction and fertility preservation.

Conflict of interest statement

Nothing declared.

Acknowledgements

The authors would like to acknowledge J.S. Del Valle and E.M. Lam for the histological images (hematoxylin and eosin staining) of human adult ovaries used in Figure 2. This work was supported by the European Research Council Consolidator Grant OVOGROWTH (ERC-CoG-2016-725722) and the Chinese Scholarship Council (CSC 201706320328).

References

Papers of particular interest, published within the period of review, have been highlighted as:

- * of special interest
- ** of outstanding interest

1. Sanchez F, Smits J: **Molecular control of oogenesis**. *Biochim Biophys Acta* 2012, **1822**:1896–1912.
2. Clarke HJ: *Chapter 2 control of Mammalian oocyte development by interactions with the maternal follicular environment*. 2017.
3. Baerwald AR, Adams GP, Pierson RA: **Ovarian antral folliculogenesis during the human menstrual cycle: a review**. *Hum Reprod Update* 2012, **18**:73–91.
4. Young JM, McNeilly AS: **Theca: the forgotten cell of the ovarian follicle**. *Reproduction* 2010, **140**:489–504.
5. Fragouli E, Lalioti MD, Wells D: **The transcriptome of follicular cells: biological insights and clinical implications for the treatment of infertility**. *Hum Reprod Update* 2014, **20**:1–11.
6. Gougeon A: **Human ovarian follicular development: from activation of resting follicles to preovulatory maturation**. *Ann Endocrinol (Paris)* 2010, **71**:132–143.
7. Devoto L, Fuentes A, Kohen P, et al.: **The human corpus luteum: life cycle and function in natural cycles**. *Fertil Steril* 2009, **92**:1067–1079.
8. Abedel-Majed MA, Romereim SM, Davis JS, Cupp AS: **Perturbations in lineage specification of granulosa and theca cells may alter corpus luteum formation and function**. *Front Endocrinol (Lausanne)* 2019, **10**:832.
9. Marozzi S, Rossi V, Salustri A, et al.: **Programmed cell death in the human ovary**. *Minerva Ginecol* 2018, **70**:549–560.
10. Fan X, Bialecka M, Moustakas I, et al.: **Single-cell reconstruction of follicular remodeling in the human adult ovary**. *Nat Commun* 2019, **10**:3164.

This article described somatic cell types and transcriptional signatures of human adult ovarian cells from medulla and antral follicles sized 1–5 mm by single-cell RNA sequencing. Several subtypes of endothelial, smooth muscle and immune cells were confirmed in human adult ovary. The analysis also revealed features of early GC and early

TC. Moreover, this paper also identified novel markers for GC and TC in antral follicles.

11. Man L, Lustgarten-Guahmich N, Kallinos E, *et al.*: **Comparison of human antral follicles of xenograft versus ovarian origin reveals disparate molecular signatures.** *Cell Rep* 2020, **32**:108027.
This article described the transcriptional profiles of GC and TC from antral follicles (1–4 mm) by single-cell RNA sequencing and included mouse-xenografted ovarian tissue. In addition, this paper applied several cell surface markers to sort GC and TC and validated some of those on tissue sections.
12. Wagner M, Yoshihara M, Douagi I, *et al.*: **Single-cell analysis of human ovarian cortex identifies distinct cell populations but no oogonial stem cells.** *Nat Commun* 2020, **11**:1147.
This article analyzed the transcriptional profile of cells from the human ovarian cortex from patients undergoing either cesarian-section or gender-reassignment surgery. The ovarian cells immunostained for DDX4 clustered with perivascular cells.
13. Zhang Y, Yan Z, Qin Q, *et al.*: **Transcriptome landscape of human folliculogenesis reveals oocyte and granulosa cell interactions.** *Mol Cell* 2018, **72**:1021–1034 e1024.
This article studied the transcriptional dynamics of oocytes and GC from follicles smaller than 300 μ m (including primordial, primary, secondary/pre-antral and early antral stages) and preovulatory MII stage. The analysis not only described transcriptional profiles of oocyte and GC from different stages, but also highlighted signaling pathways that may be involved in primordial follicle activation and oocyte-GC crosstalk.
14. Kinnear HM, Tomaszewski CE, Chang FL, *et al.*: **The ovarian stroma as a new frontier.** *Reproduction* 2020, **160**:R25–R39.
15. Arroyo A, Kim B, Yeh J: **Luteinizing hormone action in human oocyte maturation and quality: signaling pathways, regulation, and clinical impact.** *Reprod Sci* 2020, **27**:1223–1252.
16. Virant-Klun I, Knez K, Tomazevic T, Skutella T: **Gene expression profiling of human oocytes developed and matured in vivo or in vitro.** *BioMed Res Int* 2013, **2013**:879489.
17. Sanfins A, Rodrigues P, Albertini DF: **GDF-9 and BMP-15 direct the follicle symphony.** *J Assist Reprod Genet* 2018, **35**:1741–1750.
18. Markholt S, Grondahl ML, Ernst EH, *et al.*: **Global gene analysis of oocytes from early stages in human folliculogenesis shows high expression of novel genes in reproduction.** *Mol Hum Reprod* 2012, **18**:96–110.
19. Ernst EH, Grondahl ML, Grund S, *et al.*: **Dormancy and activation of human oocytes from primordial and primary follicles: molecular clues to oocyte regulation.** *Hum Reprod* 2017, **32**:1684–1700.
20. Takeuchi A, Koga K, Satake E, *et al.*: **Endometriosis triggers excessive activation of primordial follicles via PI3K-PTEN-Akt-Foxo3 pathway.** *J Clin Endocrinol Metab* 2019, **104**:5547–5554.
21. Grosbois J, Demeestere I: **Dynamics of PI3K and Hippo signaling pathways during in vitro human follicle activation.** *Hum Reprod* 2018, **33**:1705–1714.
22. Assou S, Anahory T, Pantesco V, *et al.*: **The human cumulus-oocyte complex gene-expression profile.** *Hum Reprod* 2006, **21**:1705–1719.
23. Yu B, Doni Jayavelu N, Battle SL, *et al.*: **Single-cell analysis of transcriptome and DNA methylome in human oocyte maturation.** *PLoS One* 2020, **15**, e0241698.
24. Qi L, Liu B, Chen X, *et al.*: **Single-cell transcriptomic analysis reveals mitochondrial dynamics in oocytes of patients with polycystic ovary syndrome.** *Front Genet* 2020, **11**:396.
25. Ye M, Yang ZY, Zhang Y, *et al.*: **Single-cell multiomic analysis of in vivo and in vitro matured human oocytes.** *Hum Reprod* 2020, **35**:886–900.
26. Zhao H, Li T, Zhao Y, *et al.*: **Single-cell transcriptomics of human oocytes: environment-driven metabolic competition and compensatory mechanisms during oocyte maturation.** *Antioxidants Redox Signal* 2019, **30**:542–559.
27. Ferrero H, Corachan A, Aguilar A, *et al.*: **Single-cell RNA sequencing of oocytes from ovarian endometriosis patients reveals a differential transcriptomic profile associated with lower quality.** *Hum Reprod* 2019, **34**:1302–1312.
28. Li Y, Liu H, Yu Q, *et al.*: **Growth hormone promotes in vitro maturation of human oocytes.** *Front Endocrinol (Lausanne)* 2019, **10**:485.
29. Zou H, Chen B, Ding D, *et al.*: **Melatonin promotes the development of immature oocytes from the COH cycle into healthy offspring by protecting mitochondrial function.** *J Pineal Res* 2020, **68**, e12621.
30. Li J, Lu M, Zhang P, *et al.*: **Aberrant spliceosome expression and altered alternative splicing events correlate with maturation deficiency in human oocytes.** *Cell Cycle* 2020:1–13.
31. Barone S, Sarogni P, Valli R, *et al.*: **Chromosome missegregation in single human oocytes is related to the age and gene expression profile.** *Int J Mol Sci* 2020:21.
32. Zhang JJ, Liu X, Chen L, *et al.*: **Advanced maternal age alters expression of maternal effect genes that are essential for human oocyte quality.** *Aging (Albany NY)* 2020, **12**:3950–3961.
33. Li L, Dong J, Yan L, *et al.*: **Single-cell RNA-seq analysis maps development of human germline cells and gonadal niche interactions.** *Cell Stem Cell* 2017, **20**:858–873 e854.
34. Laitinen MP, Anttonen M, Ketola I, *et al.*: **Transcription factors GATA-4 and GATA-6 and a GATA family cofactor, FOG-2, are expressed in human ovary and sex cord-derived ovarian tumors.** *J Clin Endocrinol Metab* 2000, **85**:3476–3483.
35. Ford EA, Beckett EL, Roman SD, *et al.*: **Advances in human primordial follicle activation and premature ovarian insufficiency.** *Reproduction* 2020, **159**:R15–R29.
36. Grosbois J, Devos M, Demeestere I: **Implications of non-physiological ovarian primordial follicle activation for fertility preservation.** *Endocr Rev* 2020, **41**:847–872.
37. Ernst EH, Franks S, Hardy K, *et al.*: **Granulosa cells from human primordial and primary follicles show differential global gene expression profiles.** *Hum Reprod* 2018, **33**:666–679.
38. Wyse BA, Fuchs Weizman N, Kadish S, *et al.*: **Transcriptomics of cumulus cells - a window into oocyte maturation in humans.** *J Ovarian Res* 2020, **13**:93.
39. Poulsen LC, Englund ALM, Wissing MLM, *et al.*: **Human granulosa cells function as innate immune cells executing an inflammatory reaction during ovulation: a microarray analysis.** *Mol Cell Endocrinol* 2019, **486**:34–46.
40. Poulsen LC, Botkjaer JA, Ostrup O, *et al.*: **Two waves of transcriptomic changes in periovulatory human granulosa cells.** *Hum Reprod* 2020, **35**:1230–1245.
This article analyzed the transcriptional profiles of human cumulus GC during the ovulation period, highlighting two waves of transcriptional changes. Genes related to different functions and pathways were found enriched at different time points during ovulation.
41. Poulsen LC, Pla I, Sanchez A, *et al.*: **Progressive changes in human follicular fluid composition over the course of ovulation: quantitative proteomic analyses.** *Mol Cell Endocrinol* 2019, **495**:110522.
42. Asiabi P, Leonel ECR, Marbaix E, *et al.*: **Immunodetection and quantification of enzymatic markers in theca cells: the early process of ovarian steroidogenesis.** *Biol Reprod* 2020, **102**:145–155.
43. Kamat BR, Brown LF, Manseau EJ, *et al.*: **Expression of vascular permeability factor/vascular endothelial growth factor by human granulosa and theca lutein cells. Role in corpus luteum development.** *Am J Pathol* 1995, **146**:157–165.
44. Maybin JA, Duncan WC: **The human corpus luteum: which cells have progesterone receptors?** *Reproduction* 2004, **128**:423–431.
45. Bagnjuk K, Mayerhofer A: **Human luteinized granulosa cells-A cellular model for the human corpus luteum.** *Front Endocrinol (Lausanne)* 2019, **10**:452.

46. Tajima K, Orisaka M, Mori T, Kotsuji F: **Ovarian theca cells in follicular function.** *Reprod Biomed Online* 2007, **15**:591–609.
47. Richards JS, Ren YA, Candelaria N, *et al.*: **Ovarian follicular theca cell recruitment, differentiation, and impact on fertility: 2017 update.** *Endocr Rev* 2018, **39**:1–20.
48. Groten T, Fraser HM, Duncan WC, *et al.*: **Cell junctional proteins in the human corpus luteum: changes during the normal cycle and after HCG treatment.** *Hum Reprod* 2006, **21**: 3096–3102.
49. Auersperg N, Wong AS, Choi KC, *et al.*: **Ovarian surface epithelium: biology, endocrinology, and pathology.** *Endocr Rev* 2001, **22**:255–288.
50. Ng A, Barker N: **Ovary and fimbrial stem cells: biology, niche and cancer origins.** *Nat Rev Mol Cell Biol* 2015, **16**:625–638.
51. Schindler AJ, Watanabe A, Howell SB: **LGR5 and LGR6 in stem cell biology and ovarian cancer.** *Oncotarget* 2018, **9**: 1346–1355.